

$|V_{ub}|$ MEASUREMENTS AT B -FACTORIES

A SARTI

Dep. of Physics, The University of Ferrara and INFN, I-44100 Ferrara, Italy

Abstract

The determination of the $|V_{ub}|$ element of the Cabibbo-Kobayashi-Maskawa (CKM) quark mixing matrix plays a central role in testing the Standard Model (SM) interpretation of CP violation. Measurements at B -Factories are contributing with results that rely on different theoretical assumptions. Recent inclusive and exclusive results from CLEO, *BABAR* and BELLE are reviewed.

The element $|V_{ub}|$ of the Cabibbo-Kobayashi-Maskawa (CKM) matrix [1] plays a central role in tests of the unitarity of this matrix within the CP violation mechanism as predicted by the Standard Model (SM). Precision $|V_{ub}|$ measurements, based on theoretical calculations performed at tree level, free of new physics contribution expected in loop processes, can constrain the SM predictions with boundaries that should survive any future SM extension.

Given that other main parameters of the CKM matrix are known with high precision, stringent SM tests require an uncertainty on $|V_{ub}|$ of less than 10%. The present relative error, for both inclusive and exclusive measurement techniques, is $\sim 15\%$. Measuring $|V_{ub}|$ by using inclusive charmless semileptonic B decays is a major experimental task due to the very high $\bar{B} \rightarrow X_c \ell \bar{\nu}$ background that needs to be rejected when studying $\bar{B} \rightarrow X_u \ell \bar{\nu}$ decays ($\text{BR}(\bar{B} \rightarrow X_c \ell \bar{\nu}) \sim 60 \text{ BR}(\bar{B} \rightarrow X_u \ell \bar{\nu})$). Kinematic variables describing the semileptonic decay, such as the lepton energy (E_ℓ), the invariant mass of the lepton pair (q^2) or the invariant mass of hadronic system (m_X), can be used with different power to discriminate between $b \rightarrow c$ and $b \rightarrow u$ transitions.

Theoretically, the introduction of cuts for the background suppression introduces additional errors due to the uncertainties on the parametrization of Fermi motion of

the b -quark inside the B -meson[2]. The $|V_{ub}|$ measurement using exclusive decays is also challenging from the theoretical point of view: analyses deal with many signal models and Form Factors (FF) lattice calculations. One main experimental advantage of measurements done at B factories is the possible full reconstruction of a decaying B meson that allows constraints on the missing momentum (ν reconstruction) and allows a B meson clean sample selection. Sample statistics is hence reduced, but cuts can be applied on charge conservation, ν reconstruction and lepton number, in order to clean up the sample from background events.

1 Inclusive analyses

Two independent strategies are presented, based on the study of the E_ℓ endpoint spectrum and the m_X distribution. The first one is characterized by a large semileptonic events reconstruction efficiency and low signal events acceptance ($\sim 20\%$). The latter has a lower semileptonic events reconstruction efficiency, a large signal events acceptance ($\sim 50\text{-}80\%$ of them are retained by a cut on m_X) and a still competitive signal over background ratio (S/B). Both approaches are mainly affected by the theoretical error on the extrapolation to full phase space.

The study of the E_ℓ endpoint spectrum has been performed by CLEO[3] and BABAR[4]. CLEO analysis is based on a sample of 9.1 fb^{-1} on-peak (ON) and 4.3 fb^{-1} off-peak (OFF) data. Suppression of continuum background events has been performed using a neural network. The range of lepton momentum, that has been studied and computed in the Center of Mass frame, is (p_ℓ^*) 2.2-2.6 (GeV/c). This analysis uses the continuum background subtracted p_ℓ^* spectrum, after the $B\bar{B}$ background subtraction, to compute the differential branching ratio (ΔB), that can be studied in bins of p_ℓ^* . CLEO partial branching ratio result is $\Delta B(2.2\text{-}2.6 \text{ GeV}/c) = 0.230 \pm 0.015(\text{stat}) \pm 0.035(\text{sys})$. The fraction of signal events (f_u) expected in a given bin of p_ℓ^* can be estimated by theoretical calculations making use of the $b \rightarrow s\gamma$ photon energy spectrum. Using the differential BR and f_u it is possible to compute the total $\bar{B} \rightarrow X_u \ell \bar{\nu}$ BR and extract $|V_{ub}|$ according to [5], i.e. using the equation:

$$|V_{ub}| = 0.00445 \sqrt{\left(\frac{B(b \rightarrow u\ell\bar{\nu}) \cdot 1.55 \text{ ps}}{0.002 \cdot \tau_B} \right)} \times (1.0 \pm 0.020_{\text{pert}} \pm 0.052_{1/m_b^3}). \quad (1)$$

The result obtained by CLEO is $|V_{ub}| = (4.08 \pm 0.34 \pm 0.44 \pm 0.16 \pm 0.24) \times 10^{-3}$. With a very similar approach BABAR, in the p_ℓ^* range 2.3-2.6 (GeV/c), obtains $|V_{ub}| = (4.43 \pm 0.29 \pm 0.50 \pm 0.25 \pm 0.35) \times 10^{-3}$. Quoted errors are, respectively, the experimental one (statistical plus detector systematic), the theoretical estimate of f_u , the propagation of error in the above equation and the theoretical uncertainty

on the validity of f_u determination for $\bar{B} \rightarrow X_u \ell \bar{\nu}$ events using $b \rightarrow s \gamma$ decays. The other inclusive method herein presented makes use of the invariant mass of the hadronic system (m_X) in the $\bar{B} \rightarrow X \ell \bar{\nu}$ recoil of a fully reconstructed B meson (B_{reco}) to separate the $b \rightarrow u$ and the $b \rightarrow c$ transition. *BABAR* analysis [6] reconstructs a large sample of B mesons by selecting hadronic decays $B_{reco} \rightarrow D^{(*)}X$. The kinematic consistency of B_{reco} candidates is checked with two variables, the beam energy-substituted mass $m_{ES} = \sqrt{s/4 - \vec{p}_B^2}$ and the energy difference $\Delta E = E_B - \sqrt{s}/2$. Here $\sqrt{s}$ is the total energy in the $\Upsilon(4S)$ center of mass frame, and $\vec{p}_B$ and E_B denote the momentum and energy of the B_{reco} candidate in the same frame. $\Delta E = 0$ is required within three standard deviations as measured for each mode.

On the recoil side, the event selection proceeds via a lepton selection with a cut on the lepton momentum ($\sim 1 \text{ GeV}/c$) and on the missing mass of the event. Neutral and charged kaons are vetoed in the signal region and events with a kaon positively identified in the recoil are used as a control sample.

In order to reduce systematic uncertainties, the ratio of branching ratios $R_{u/sl} = \mathcal{B}(\bar{B} \rightarrow X_u \ell \bar{\nu})/\mathcal{B}(\bar{B} \rightarrow X \ell \bar{\nu})$ is determined from N_u , the observed number of $\bar{B} \rightarrow X_u \ell \bar{\nu}$ candidates with $m_X < 1.55 \text{ GeV}/c^2$, and N_{sl} , the number of events with at least one charged lepton:

$$R_{u/sl} = \frac{\mathcal{B}(\bar{B} \rightarrow X_u \ell \bar{\nu})}{\mathcal{B}(\bar{B} \rightarrow X \ell \bar{\nu})} = \frac{N_u/(\varepsilon_{sel}^u \varepsilon_{m_X}^u)}{N_{sl}} \times \frac{\varepsilon_l^{sl} \varepsilon_{reco}^{sl}}{\varepsilon_l^u \varepsilon_{reco}^u}. \quad (2)$$

Here ε_{sel}^u is the efficiency for selecting $\bar{B} \rightarrow X_u \ell \bar{\nu}$ decays once a $\bar{B} \rightarrow X \ell \bar{\nu}$ candidate has been identified; $\varepsilon_{m_X}^u$ is the fraction of signal events with $m_X < 1.55 \text{ GeV}/c^2$; $\varepsilon_l^{sl}/\varepsilon_l^u$ corrects for the difference in the efficiency of the lepton momentum cut for $\bar{B} \rightarrow X \ell \bar{\nu}$ and $\bar{B} \rightarrow X_u \ell \bar{\nu}$ decays, and $\varepsilon_{reco}^{sl}/\varepsilon_{reco}^u$ accounts for a possible efficiency difference in the B_{reco} reconstruction in events with $\bar{B} \rightarrow X \ell \bar{\nu}$ and $\bar{B} \rightarrow X_u \ell \bar{\nu}$ decays. N_{sl} is derived from a fit to the m_{ES} distribution. N_u is extracted from the m_X distribution by a minimum χ^2 fit to the sum of three contributions: the signal, the background N_c from $\bar{B} \rightarrow X_c \ell \bar{\nu}$, and a background of $< 1\%$ from other sources (misidentified leptons, secondary τ and charm decays). Fig. 1a shows the fitted m_X distribution. Fig. 1b shows the m_X distribution after background subtraction. By using 82 fb^{-1} integrated luminosity on the $\Upsilon(4S)$ peak, *BABAR* obtains $|V_{ub}| = (4.62 \pm 0.28 \pm 0.27 \pm 0.40 \pm 0.26) \times 10^{-3}$, where errors are statistical, detector systematic, theoretical model and propagation of error in eq.1. The S/B ratio is 1.7 (higher than any previous inclusive analysis) and the main error comes from the Fermi motion parametrization. In figure 2, left plot, inclusive results for $|V_{ub}|$ measurement are summarized together with previous results from LEP experiments.




Figure 1: The m_X distribution for $\bar{B} \rightarrow X \ell \bar{\nu}$ candidates for *BABAR*[6] inclusive analysis: a) data (points) and fit components, and b) data and signal MC after subtraction of the $b \rightarrow c \ell \nu$ and the “other” backgrounds.

2 Exclusive analyses

BABAR[9], *CLEO*[10] and *BELLE*[11] exclusive analysis strategies, based on the study of $B \rightarrow (\pi, \rho, \omega) \ell \nu$ decays, are outlined below. The signal events selection is done using the distributions of the invariant mass of $\pi\pi$ or $\pi\pi\pi$ system, the difference between the expected and reconstructed B meson energy (ΔE) and lepton momentum (p_ℓ), for $B \rightarrow n\pi \ell \nu$ decays. *CLEO* is using, for the $\rho(\omega)$ decay mode, a simultaneous fit to $m_{\pi\pi}(m_{\pi\pi\pi})$ and ΔE distributions.

The *CLEO* analysis achieves a low signal and detector modeling dependence by performing the full differential analysis in bins of q^2 . The lepton momentum ranges covered are $p_\ell > 1.0$ GeV/c (pseudo-scalar decay mode) and $p_\ell > 1.5$ GeV/c (vector decay mode). The main sources of systematics are: the modelization of K_L^0 energy deposit, the estimate of tracking efficiency, the simulation of charged and neutral particles interactions in the electromagnetic calorimeter and the physics model used to convert BR into $|V_{ub}|$. On the simulation side, signal modeling (various Form Factors for different exclusive decays) and non-resonant contributions are playing a central role. The efficiency for semileptonic and signal events selection, the continuum rejection and cross-feed background rates are depending on the event q^2 , therefore possible bias can be introduced by cuts on this variable and should be taken into account. A differential analysis that studies $d\Gamma/dq^2$ minimizes bias effects and allows to test different theoretical models. Fig. 2, right plot, shows results for $B^0 \rightarrow \pi^- \ell^+ \nu$ decays: data (markers) and the best fit to the predicted $d\Gamma/dq^2$ (histograms) for the three models used to extract both rates and $|V_{ub}|$ are




Figure 2: Left: $|V_{ub}|$ results from inclusive analyses. LEP and B factories results are shown. Right: Measured branching fractions for $B^0 \rightarrow \pi^- \ell^+ \nu$ (points) and the best fit to the predicted $d\Gamma/dq^2$ (histograms) for the three models used to extract both rates and $|V_{ub}|$. The data points have small horizontal offsets introduced for clarity.

shown. By improving precision it will be possible to treat data with more appropriate theoretical models (for example in fig. 2 the ISGWII model seems disfavored). *BABAR* and *CLEO* have followed similar approaches analyzing different p_ℓ ranges ($p_\ell > 2.3 \text{ GeV}/c$ for *BABAR* and $p_\ell > 2.4 \text{ GeV}/c$ for *BELLE*). Table 1 shows the results for all the various exclusive analyses. Attention must be paid when making averages: *CLEO* '03 analyses, for example, are using different models for $|V_{ub}|$ extraction with respect to the other analyses and it is not clear how to take it properly into account when averaging.

Analysis	Result
CLEO (ρ)	$3.23 \pm 0.24^{+0.23}_{-0.26} \pm 0.58$
BABAR (ρ)	$3.64 \pm 0.22 \pm 0.25^{+0.39}_{-0.56}$
BELLE (π)	$3.11 \pm 0.13 \pm 0.24 \pm 0.061$
BELLE (ρ^0)	$3.50 \pm 0.20 \pm 0.28$
CLEO '03 (π)	$3.24 \pm 0.22 \pm 0.13 \pm 0.09^{+0.55}_{-0.39}$
CLEO '03 (ρ)	$3.00 \pm 0.21^{+0.29}_{-0.35} \pm 0.28^{+0.49}_{-0.38}$
CLEO '03 (comb)	$3.17 \pm 0.17^{+0.16}_{-0.17} \pm 0.03^{+0.53}_{-0.39}$

Table 1: Exclusive results for $|V_{ub}|$. Quoted errors are statistical, experimental systematic, theoretical, and signal Form Factor shape, respectively.

3 Conclusions and outlook

Several new results concerning the inclusive and exclusive determination of $|V_{ub}|$ CKM matrix element have been obtained recently. The B-factory era has introduced new experimental methods based on studies of the q^2 and mass spectrum in the recoil of fully reconstructed B mesons. Together with an impressive theoretical progress (HQET and OPE for inclusive decays, FF determination for the exclusive ones) these improvements have lead to a 13% precision measurement (inclusive) and at a first attempt of discriminating theoretical models via differential analysis (exclusive). The main open issues for inclusive analyses are the ν reconstruction and the theoretical error reduction, while the exclusive ones are still suffering from a poor FF determination. A $|V_{ub}|$ measurement with an error less than 10% by 2007, and a consistency test for exclusive and inclusive determinations of $|V_{ub}|$, the proper way to combine results being still under discussion, are the main goals regarding $|V_{ub}|$ measurements in the near future.

References

- [1] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Th. Phys. **49**, 652 (1973).
- [2] F. De Fazio and M. Neubert, JHEP **9906**, 017 (1999).
- [3] A. Bornheim *et al.* [CLEO Collaboration], Phys. Rev. Lett. **88**:231803, 2002.
- [4] B. Aubert *et al.* [BABAR Collaboration], arXiv:hep-ex/0207081
- [5] K. Hagiwara *et al.* [Particle Data Group Collaboration], Phys. Rev. D **66**, 010001 (2002).
- [6] D. del Re [BABAR Collaboration], arXiv:hep-ex/0306026.
- [7] A. Sugiyama [BELLE Collaboration], arXiv:hep-ex/0306020.
- [8] A. Bornheim *et al.* [CLEO Collaboration], arXiv:hep-ex/0207064
- [9] B. Aubert *et al.* [BABAR Collaboration], e-Print Archive: hep-ex/0301001
- [10] A. Bornheim [CLEO Collaboration], arXiv:hep-ex/0307011.
- [11] M. Yamauchi [Belle Collaboration], Nucl. Phys. Proc. Suppl. **117** (2003) 83.
K. Abe *et al.* [Belle Collaboration], arXiv:hep-ex/0307075.